

BOUNDED COHOMOLOGY, CROSS RATIOS AND COCYCLES

URSULA HAMENSTÄDT

ABSTRACT. We use cross ratios to describe second real continuous bounded cohomology for locally compact σ -compact topological groups. We also investigate the second continuous bounded cohomology group of a closed subgroup of the isometry group $\text{Iso}(X)$ of a proper hyperbolic geodesic metric space X and derive some rigidity results for $\text{Iso}(X)$ -valued cocycles.

1. INTRODUCTION

A geodesic metric space X is called δ -hyperbolic for some $\delta > 0$ if it satisfies the δ -thin triangle condition: For every geodesic triangle in X with sides a, b, c the side a is contained in the δ -neighborhood of $b \cup c$. If X is proper then X can naturally be compactified by adding the *Gromov boundary* ∂X . The isometry group $\text{Iso}(X)$ of X , equipped with the compact open topology, is a locally compact σ -compact topological group which acts on $X \cup \partial X$ as a group of homeomorphisms. The *limit set* of a closed subgroup G of $\text{Iso}(X)$ is the set of accumulation points in ∂X of an orbit of the action of G on X ; this limit set is a compact G -invariant subset of ∂X . The group G is called *elementary* if its limit set consists of at most two points.

Let S be a standard Borel space and let μ be a Borel probability measure on S . Let Γ be a countable group which admits a measure preserving ergodic action on (S, μ) . An $\text{Iso}(X)$ -valued *cocycle* for this action is a measurable map $\alpha : \Gamma \times S \rightarrow \text{Iso}(X)$ such that $\alpha(gh, x) = \alpha(g, hx)\alpha(h, x)$ for all $g, h \in \Gamma$ and μ -almost every $x \in S$. The cocycle α is *cohomologous* to a cocycle $\beta : \Gamma \times S \rightarrow \text{Iso}(X)$ if there is a measurable map $\varphi : S \rightarrow \text{Iso}(X)$ such that $\varphi(gx)\alpha(g, x) = \beta(g, x)\varphi(x)$ for all $g \in G$, μ -almost every $x \in S$. Recall that a *compact extension* of a locally compact topological group G is a locally compact topological group H which contains a normal compact subgroup K such that $G = H/K$ as topological groups. Extending earlier results of Monod and Shalom [14] we show in Section 4.

Theorem A: *Let G be a semi-simple Lie group with finite center, no compact factors and of rank at least 2. Let $\Gamma < G$ be an irreducible lattice which admits a mixing measure preserving action on a probability space (S, μ) . Let X be a proper hyperbolic geodesic metric space and let $\alpha : \Gamma \times S \rightarrow \text{Iso}(X)$ be a cocycle; then one of the following two possibilities holds.*

- (1) α is cohomologous to a cocycle into an elementary subgroup of $\text{Iso}(X)$.

Date: August 24, 2005.

Partially supported by Sonderforschungsbereich 611.

- (2) α is cohomologous to a cocycle into a subgroup H of $\text{Iso}(X)$ which is a compact extension of a simple Lie group L of rank one, and there is a continuous surjective homomorphism $G \rightarrow L$.

The proof of this result uses second bounded cohomology for closed subgroups of the isometry group $\text{Iso}(X)$ of X . Here the second bounded cohomology group of a locally compact topological group G with coefficients in a *Banach module* E for G , i.e. a Banach space E together with a continuous homomorphism of G into the group of linear isometries of E , is defined as follows. For every $i \geq 1$, the group G naturally acts on the vector space $C_b(G^i, E)$ of continuous bounded maps $G^i \rightarrow E$. If we denote by $C_b(G^i, E)^G \subset C_b(G^i, E)$ the linear subspace of all G -invariant such maps, then the *second continuous bounded cohomology group* $H_{cb}^2(G, E)$ of G with coefficients E is defined as the second cohomology group of the complex

$$(1) \quad 0 \rightarrow C_b(G, E)^G \xrightarrow{d} C_b(G^2, E)^G \xrightarrow{d} \dots$$

with the usual homogeneous coboundary operator d (see [13]). If G is a countable group with the discrete topology then we write $H_b^2(G, E) = H_{cb}^2(G, E)$. Of particular importance is the case that $E = \mathbb{R}$ with the trivial G -action which yields real continuous second bounded cohomology. There is a natural map of $H_{cb}^2(G, \mathbb{R})$ to the ordinary continuous second cohomology group $H_c^2(G, \mathbb{R})$ of G which in general is neither injective nor surjective.

If G is a non-elementary closed subgroup of the isometry group of a proper hyperbolic geodesic metric space X with limit set $\Lambda \subset \partial X$ which does not act transitively on the complement of the diagonal in $\Lambda \times \Lambda$ then the kernel of the natural map $H_{cb}^2(G, \mathbb{R}) \rightarrow H_c^2(G, \mathbb{R})$ is infinite dimensional [9]. In contrast, we show in Section 3.

Theorem B: *Let X be a proper hyperbolic geodesic metric space and let $G < \text{Iso}(X)$ be a non-elementary closed subgroup with limit set Λ . If G acts transitively on the complement of the diagonal in $\Lambda \times \Lambda$ then the kernel of the natural map $H_{cb}^2(G, \mathbb{R}) \rightarrow H_c^2(G, \mathbb{R})$ is trivial.*

In Section 2 we identify for an arbitrary locally compact σ -compact topological group G the group $H_{cb}^2(G, \mathbb{R})$ with the vector space of measurable anti-symmetric G -cross ratios on a strong boundary for G . As an application we describe the kernel of the natural map $H_b^2(\Gamma, \mathbb{R}) \rightarrow H^2(\Gamma, \mathbb{R})$ for a lattice Γ in $PSL(2, \mathbb{R})$. Namely, recall that Γ acts as a group of orientation preserving isometries on the hyperbolic plane $\mathbf{H}^2$ and hence it acts on the boundary $S^1 = \partial \mathbf{H}^2$ of $\mathbf{H}^2$ preserving the measure class of the Lebesgue measure λ . We show that the kernel of the natural map $H_b^2(\Gamma, \mathbb{R}) \rightarrow H^2(\Gamma, \mathbb{R})$ is naturally isomorphic to the vector space of all finite Γ -invariant reflection-anti-invariant finitely additive signed measures μ on the complement of the diagonal in $S^1 \times S^1$ with the additional property that the map $(a, b, c, d) \in S^1 \rightarrow \mu[a, b) \times [c, d)$ is λ -measurable.

2. CROSS-RATIOS AND BOUNDED COHOMOLOGY

Let X be an arbitrary set with infinitely many points. We define an *anti-symmetric cross ratio* for X to be a bounded function $[\cdot]$ on the space of quadruples of pairwise distinct points in X with the following additional properties.

- i) $[\xi, \xi', \eta, \eta'] = -[\xi', \xi, \eta, \eta']$.
- ii) $[\eta, \eta', \xi, \xi'] = -[\xi, \xi', \eta, \eta']$.
- iii) $[\xi, \xi', \eta, \eta'] + [\xi', \xi'', \eta, \eta'] = [\xi, \xi'', \eta, \eta']$.

Note that $[\cdot]$ can naturally be extended by 0 to the set of quadruples of points in X of the form (ξ, ξ, η, ζ) where ξ, η, ζ are pairwise distinct.

Let Γ be an arbitrary countable group which acts on X . We define a Γ -*cross ratio* on X to be an anti-symmetric cross ratio on X which is invariant under the action of Γ . Let $\mathcal{C}(\Gamma, X)$ be the vector space of all anti-symmetric Γ -cross ratios on X , equipped with the supremums-norm $\|\cdot\|$. Thus we have $\|[\cdot]\| = \sup\{[\xi, \xi', \eta, \eta'] \mid \xi, \xi', \eta, \eta' \in X \text{ are pairwise distinct}\}$, and $(\mathcal{C}(\Gamma, X), \|\cdot\|)$ is naturally a Banach space.

The second real bounded cohomology group $H_b^2(\Gamma, \mathbb{R})$ of Γ is a vector space of equivalence classes of bounded functions on $\Gamma \times \Gamma \times \Gamma$ which are invariant under the free left action of Γ . It can be equipped with a natural pseudo-norm which assigns to a cohomology class the infimum of the supremums-norms of any representative of the class. For the second bounded cohomology, this pseudo-norm turns out to be a norm which provides $H_b^2(\Gamma, \mathbb{R})$ with the structure of a Banach space. We have.

Proposition 2.1: *There is a continuous linear map $\Omega : \mathcal{C}(\Gamma, X) \rightarrow H_b^2(\Gamma, \mathbb{R})$.*

Proof: Let $[\cdot] \in \mathcal{C}(\Gamma, X)$ and let (ξ, ζ, η) be a triple of pairwise distinct points in X . Choose an arbitrary point $\nu \in X$ which is distinct from any of the points in the triple and define

$$(2) \quad \varphi(\xi, \eta, \zeta) = \frac{1}{2}([\xi, \zeta, \eta, \nu] + [\zeta, \eta, \xi, \nu] + [\eta, \xi, \zeta, \nu]).$$

We claim first that this does not depend on the choice of ν . Namely, let $\nu' \in X - \{\xi, \eta, \zeta\}$ be arbitrary. We compute

$$(3) \quad \begin{aligned} & [\xi, \zeta, \eta, \nu] + [\zeta, \eta, \xi, \nu] + [\eta, \xi, \zeta, \nu] - [\xi, \zeta, \eta, \nu'] - [\zeta, \eta, \xi, \nu'] - [\eta, \xi, \zeta, \nu'] \\ &= [\xi, \zeta, \nu', \eta] + [\xi, \zeta, \eta, \nu] + [\zeta, \eta, \nu', \xi] + [\zeta, \eta, \xi, \nu] + [\eta, \xi, \nu', \zeta] + [\eta, \xi, \zeta, \nu] \\ &= [\xi, \zeta, \nu', \nu] + [\zeta, \eta, \nu', \nu] + [\eta, \xi, \nu', \nu] = 0 \end{aligned}$$

by the defining properties of an anti-symmetric cross ratio. This shows our claim. As a consequence, the function φ is well defined, moreover it is invariant under the action of the group Γ . Its supremums norm is bounded in absolute value by $3\|[\cdot]\|/2$. We extend φ by 0 to the set of all triples of points in X for which at least two of the points coincide. Then φ induces a bounded function μ on Γ^3 which is invariant under the free left action of Γ as follows. Choose any point $\xi \in X$ and define $\mu(\gamma_1, \gamma_2, \gamma_3) = \varphi(\gamma_1\xi, \gamma_2\xi, \gamma_3\xi)$.

We next show that this function μ is a cocycle, i.e. that it is alternating and its image under the natural coboundary map $L^\infty(\Gamma^3, \mathbb{R}) \rightarrow L^\infty(\Gamma^4, \mathbb{R})$ vanishes. For this it is enough to show the corresponding properties for the function φ . Namely, observe that for a triple (a, b, c) of pairwise distinct points in X and any $d \in X - \{a, b, c\}$ we have

$$(4) \quad \begin{aligned} 2(\varphi(a, c, b) + \varphi(a, b, c)) &= [a, b, c, d] + [b, c, a, d] + [c, a, b, d] \\ &+ [a, c, b, d] + [c, b, a, d] + [b, a, c, d] = 0 = 2(\varphi(a, c, b) + \varphi(c, a, b)). \end{aligned}$$

which shows that φ is alternating. Next we have to show that

$$(5) \quad \varphi(b, c, d) - \varphi(a, c, d) + \varphi(a, b, d) - \varphi(a, b, c) = 0$$

for any quadruple (a, b, c, d) of points in X . By our observation (4), equation (5) is valid for every quadruple (a, b, c, d) of points in X for which at least 2 of the points coincide. On the other hand, for every quadruple (a, b, c, d) of pairwise distinct points in X we have

$$(6) \quad \begin{aligned} 2(\varphi(a, b, c) - \varphi(b, c, d)) &= [a, c, b, d] + [c, b, a, d] + [b, a, c, d] \\ &- [b, d, c, a] - [d, c, b, a] - [c, b, d, a] = 2[c, b, a, d] \end{aligned}$$

by the defining properties of an anti-symmetric cross ratio. Together with equation (4) we deduce with the same calculation as before that $\varphi(a, c, d) - \varphi(a, b, d) = -\varphi(c, a, d) + \varphi(a, d, b) = -[d, a, c, b] = -[c, b, a, d]$ and consequently φ satisfies the cocycle equation (5). Therefore the function μ on Γ defines a bounded cocycle and hence a second bounded cohomology class $\Omega([\cdot]) \in H_b^2(\Gamma, \mathbb{R})$.

We next observe that the cohomology class $\Omega([\cdot])$ does not depend on the choice of the point $\xi \in X$ which we used to define the function φ . Namely, let η be a different point in X and define $\mu'(\gamma_1, \gamma_2, \gamma_3) = \varphi(\gamma_1\eta, \gamma_2\eta, \gamma_3\eta)$. For $\gamma \in \Gamma$ write $\nu(\gamma) = [\gamma\xi, \eta, \xi, \gamma\eta]$. By equation (4), (6) we have $\nu(\gamma) = \varphi(\xi, \eta, \gamma\xi) + \varphi(\gamma\xi, \eta, \gamma\eta)$ and consequently $\mu(e, \gamma_1, \gamma_2) - \mu'(e, \gamma_1, \gamma_2) - \nu(\gamma_1) - \nu(\gamma_1^{-1}\gamma_2) + \nu(\gamma_2)$ is the value of φ on the boundary of a singular polyhedron of dimension 3 whose vertices consist of the points $\xi, \eta, \gamma_1\xi, \gamma_1\eta, \gamma_2\xi, \gamma_2\eta$ and whose sides contain the simplices with vertices $\xi, \gamma_1\xi, \gamma_2\xi$ and $\eta, \gamma_1\eta, \gamma_2\eta$ as well as three quadrangles with the same set of vertices. Since φ is a cocycle, the evaluation of φ on this polyhedron vanishes and therefore the function $\mu - \mu'$ is the image of a Γ -invariant bounded function on Γ^2 under the coboundary map $L^\infty(\Gamma^2) \rightarrow L^\infty(\Gamma^3)$. As a consequence, for every $[\cdot] \in \mathcal{C}(\Gamma, X)$ the bounded cohomology class $\Omega([\cdot]) \in H_b^2(\Gamma, \mathbb{R})$ only depends on the cross ratio $[\cdot]$ but not on the choice of a point $\xi \in X$. The resulting map $\Omega : \mathcal{C}(\Gamma, X) \rightarrow H_b^2(\Gamma, \mathbb{R})$ is clearly linear and continuous with respect to the supremum norm on $\mathcal{C}(\Gamma, X)$ and the Gromov norm on $H_b^2(\Gamma, \mathbb{R})$. $\square$

For a topological space X define a *continuous* anti-symmetric cross ratio to be a continuous function $[\cdot]$ on the space of quadruples of pairwise distinct points in X which is an anti-symmetric cross ratio in the sense of our definition. If G is any locally compact σ -compact topological group which acts on a locally compact σ -compact topological space X as a group of homeomorphisms, then we define a continuous anti-symmetric G -cross-ratio to be a continuous anti-symmetric cross ratio on X which is invariant under the diagonal action of G .

Every locally compact σ -compact group G acts freely on itself by left translations. Thus we can use Proposition 2.1 for $X = G$ to deduce that there is a continuous linear map from the vector space of all continuous anti-symmetric G -cross ratios on G into $H_{cb}^2(G, \mathbb{R})$. Namely, we have.

Corollary 2.2: *Let G be a locally compact σ -compact group; then $H_{cb}^2(G, \mathbb{R})$ is the quotient of the space of continuous anti-symmetric G -cross ratios on G under the coboundary relation.*

Proof: By Proposition 2.1 and its proof there is a linear map Ω from the space of all continuous anti-symmetric G -cross ratios on G into $H_{cb}^2(G, \mathbb{R})$. We have to show that Ω is surjective. For this recall first that every second continuous bounded cohomology class for G is the class of a G -invariant continuous *alternating* function φ on G^3 which satisfies the cocycle equation (5) (see [10, 13]). For such a function φ define $[g, g', h, h'] = \varphi(h, g', g) - \varphi(g', g, h')$ as in the proof of Proposition 2.1. Then $[\]$ is continuous and G -invariant; we claim that it satisfies the requirements of an anti-symmetric G -cross ratio.

Namely, first of all we have

$$(7) \quad [b, c, a, d] = \varphi(a, c, b) - \varphi(c, b, d) = -\varphi(a, b, c) + \varphi(b, c, d) = -[c, b, a, d]$$

since φ is alternating and similarly

$$(8) \quad [a, d, c, b] = \varphi(c, d, a) - \varphi(d, a, b) = -\varphi(a, b, d) + \varphi(a, c, d) = -[c, b, a, d].$$

In the same way we obtain from the cocycle equation (5) for φ that

$$(9) \quad [a, b, c, d] + [b, b', c, d] = [a, b', c, d]$$

which shows that $[\]$ is indeed a G -invariant anti-symmetric cross ratio. By the proof of Proposition 2.1, the cohomology class of φ is just $\Omega([\])$. Thus the map Ω is surjective. $\square$

A locally compact σ -compact topological group G admits a *strong boundary* [11] which is a standard measure space (B, μ) together with a measure class preserving action of G such that the following two conditions are satisfied.

- (1) The G -action on B is amenable.
- (2) For any separable Banach- G -module E , any measurable G -equivariant map $B \times B \rightarrow E$ is essentially constant.

Define a *measurable anti-symmetric cross ratio* on B to be a *measurable* essentially bounded G -invariant function on the space of quadruples of distinct points in B which satisfies almost everywhere the defining equation for a cross ratio. Our above considerations immediately imply.

Lemma 2.3: *Let (B, μ) be a strong boundary for G ; then $H_{cb}^2(G, \mathbb{R})$ is naturally isomorphic to the space of G -invariant μ -measurable anti-symmetric cross ratios on B .*

Proof: Let G be a locally compact σ -compact group and let (B, μ) be a strong boundary for G . Denote for $k \geq 1$ by $L^\infty(B^k, \mu^k)$ the Banach space of essentially bounded μ^k -measurable functions on B^k ; it contains the closed subspace $L^\infty(B^k, \mu^k)^G$ of all G -invariant such functions. By the results of Burger and Monod [6], $H_{cb}^2(G, \mathbb{R})$ coincides with the second bounded cohomology group of the resolution

$$(10) \quad \mathbb{R} \rightarrow L^\infty(B, \mu)^G \rightarrow L^\infty(B^2, \mu^2)^G \rightarrow \dots$$

with the usual homogeneous coboundary operator [13]. By Proposition 2.1 and its proof, there is a natural bijection between the space of μ^4 -measurable essentially bounded anti-symmetric G -cross ratios on B and the space of μ^3 -measurable bounded cocycles on B , i.e. the space of alternating functions in $L^\infty(B^3, \mu^3)^G$ which satisfy the cocycle equation. Any two such bounded cocycles are cohomologous if and only if they differ by the image under the coboundary operator of an element of $L^\infty(B^2, \mu^2)^G$. However, by ergodicity of the measure class μ^2 under the diagonal action of Γ , the vector space $L^\infty(B^2, \mu^2)^G$ only contains essentially constant functions. This means that the natural map which assigns to a μ^4 -measurable G -cross ratio its corresponding bounded cohomology class is injective. $\square$

We next look at a specific example. Namely, the ideal boundary of the oriented hyperbolic plane $\mathbf{H}^2$ can naturally be identified with the oriented circle S^1 . The action on S^1 of the group $PSL(2, \mathbb{R})$ of orientation preserving isometries of $\mathbf{H}^2$ preserves the measure class of the Lebesgue measure λ . Moreover, (S^1, λ) is a strong boundary for $PSL(2, \mathbb{R})$. A triple (a, b, c) of pairwise distinct points in S^1 will be called *ordered* if the point b is contained in the interior of the oriented subinterval (a, c) of S^1 . Similarly we define a quadruple (a, b, c, d) of pairwise distinct points on S^1 to be ordered. Ordered quadruples define a $PSL(2, \mathbb{R})$ -invariant subset of the space of quadruples of pairwise distinct points in S^1 . There is a unique antisymmetric cross ratio $[\]_0$ on S^1 with the following properties.

- (1) $[a, b, c, d]_0 = 0$ if (a, b, c, d) is ordered.
- (2) $[a, b, c, d]_0 = 1$ if (a, c, b, d) is ordered.

Clearly this cross ratio is measurable and invariant under the action of $PSL(2, \mathbb{R})$. We have.

Lemma 2.4:

- (1) Up to a constant, $[\]_0$ is the unique non-trivial antisymmetric cross ratio on S^1 which is invariant under the full group $PSL(2, \mathbb{R})$ of orientation preserving isometries of $\mathbf{H}^2$.
- (2) If μ is any Borel measure on S^1 without atoms and if $[\]$ is any μ -measurable antisymmetric cross ratio on S^1 which satisfies $[a, b, c, d] = 0$ for μ -almost every ordered quadruple of pairwise distinct points in S^1 then $[\]$ is a multiple of $[\]_0$.

Proof: If $[\]$ is any antisymmetric cross ratio which is invariant under the full group $PSL(2, \mathbb{R})$ then the same is true for the induced function φ on the set of triples of pairwise distinct points in S^1 . Recall that the group $PSL(2, \mathbb{R})$ acts

transitively on the space of *ordered* triples of distinct points in S^1 and consequently we have $\varphi(a, b, c) = \varphi(b, c, d)$ for every ordered quadruple (a, b, c, d) of pairwise distinct points in S^1 . Hence equation (6) in the proof of Proposition 2.1 implies that $[a, b, c, d] = 0$ whenever the quadruple (a, b, c, d) of pairwise distinct points in S^1 is ordered. As a consequence, if $[\] \neq 0$ then after possibly multiplying $[\]$ with a constant we conclude from the definition of an anti-symmetric cross ratio that $(a, b, c, d) = 1$ whenever (a, c, b, d) is ordered. In other words, up to a constant we have $[\] = [\]_0$ which shows the first part of the lemma.

To show the second part of the lemma, let μ be a Borel measure on S^1 without atoms and let $[\]$ be any μ -measurable antisymmetric cross ratio on S^1 such that $[a, b, c, d] = 0$ for almost every ordered quadruple of pairwise distinct points in S^1 . Let (a, b, c, d) be a quadruple of pairwise distinct points in S^1 such that $[a, b, c, d] \neq 0$. By the symmetry relation of an antisymmetric cross ratio we may assume that (a, c, b, d) is ordered. Thus the point b is contained in the oriented open subinterval (c, d) of S^1 . Now if $b' \in (c, d)$ is another such point then by the defining properties of an antisymmetric cross ratio we have $[a, b, c, d] + [b, b', c, d] = [a, b', c, d]$. Assume first that $b' \in (b, d)$; then the quadruple (b, b', d, c) is ordered and consequently we have $[b, b', c, d] = 0$ by assumption and $[a, b, c, d] = [a, b', c, d]$. Via exchanging b and b' we conclude that $[a, b, c, d]$ does not depend on $b \in (c, d)$. Similarly we deduce that in fact $[a, b, c, d]$ only depends on the order of the points in the quadruple (a, b, c, d) . In other words, $[\]$ is a multiple of $[\]_0$. $\square$

Lemma 2.4 shows in particular that $[\]_0$ corresponds to a generator α of the group $H_{cb}^2(PSL(2, \mathbb{R}), \mathbb{R})$.

Now let $\Gamma < PSL(2, \mathbb{R})$ be any finitely generated torsion free discrete group. Then Γ acts freely on the hyperbolic plane $\mathbf{H}^2$ as a properly discontinuous group of isometries, and $S = \mathbf{H}^2/\Gamma$ is an oriented surface. The unit circle $S^1 = \partial\mathbf{H}^2$ with the Γ -invariant measure class of the Lebesgue measure λ is a strong boundary for Γ . If Γ is cocompact then we have $H^2(\Gamma, \mathbb{R}) = H^2(M, \mathbb{R}) = \mathbb{R}$ and there is a natural surjective restriction homomorphism $H_b^2(\Gamma, \mathbb{R}) \rightarrow H^2(\Gamma, \mathbb{R})$. The diagram

$$\begin{array}{ccc} H_b^2(\Gamma, \mathbb{R}) & \longrightarrow & H^2(\Gamma, \mathbb{R}) \\ \uparrow & & \uparrow \\ H_{cb}^2(PSL(2, \mathbb{R}), \mathbb{R}) & \longrightarrow & H_c^2(PSL(2, \mathbb{R}), \mathbb{R}) \end{array}$$

commutes and hence in our identification of $H_b^2(\Gamma, \mathbb{R})$ with the space of Γ -invariant antisymmetric cross ratios on the unit circle S^1 , the $PSL(2, \mathbb{R})$ -invariant cross ratio $[\]_0$ from Lemma 2.4 maps to a multiple of the Euler class of the tangent bundle of S . An easy calculation shows that the multiplication factor is in fact 1 (compare [4] for a detailed explanation of this fact). If Γ is not cocompact then S is homeomorphic to the interior of a compact oriented surface $\hat{S}$ whose boundary $\partial\hat{S}$ is a finite union of circles. In particular, the fundamental group of each of the boundary circles is amenable. Now the bounded cohomology of any countable cellular space X is naturally isomorphic to the bounded cohomology of its fundamental group $\pi_1(X)$ [10]. Thus $H_b(\partial\hat{S}, \mathbb{R}) = \{0\}$ and hence the relative group $H_b^2(\hat{S}, \partial\hat{S}, \mathbb{R})$ is isomorphic to $H_b^2(\hat{S}, \mathbb{R}) = H_b^2(\Gamma, \mathbb{R})$. The image under the

natural map $\iota : H_b^2(\hat{S}, \partial\hat{S}, \mathbb{R}) \rightarrow H^2(\hat{S}, \partial\hat{S}, \mathbb{R})$ of the pull-back $i^*\alpha$ of α under the inclusion $i : \Gamma \rightarrow PSL(2, \mathbb{R})$ is just the Euler class of the (bordered) surface $\hat{S}$. In other words, the evaluation of $\iota(i^*\alpha)$ on the fundamental class of the pair $(\hat{S}, \partial\hat{S})$ just equals the Euler characteristic of S (we refer to [4] for a detailed discussion of a slightly different viewpoint on these facts).

To describe the kernel Q of the natural map $H_b^2(\Gamma, \mathbb{R}) \rightarrow H^2(\hat{S}, \partial\hat{S}, \mathbb{R})$, let Δ be the diagonal in $S^1 \times S^1$ and recall that a finitely additive signed measure on $S^1 \times S^1 - \Delta$ is a function μ which assigns to each oriented rectangle of the form $[a, b] \times [c, d]$ where a, b, c, d are pairwise distinct a number $\mu[a, b] \times [c, d] \in \mathbb{R}$ such that

$$(11) \quad \mu[a, b] \times [c, d] + \mu[b, b'] \times [c, d] = \mu[a, b'] \times [c, d]$$

whenever $[a, b'] = [a, b] \cup [b, b']$ is disjoint from $[c, d]$. This measure is flip-antiinvariant if $\mu[a, b] \times [c, d] = -\mu[c, d] \times [a, b]$. Let λ be the Lebesgue measure on S^1 which defines the unique $PSL(2, \mathbb{R})$ -invariant measure class. The collection of all finitely additive flip-antiinvariant signed measures μ on $S^1 \times S^1$ with the additional property that the function $(a, b, c, d) \rightarrow \mu[a, b] \times [c, d]$ is λ -measurable is clearly a vector space; we call it the space of λ -measurable flip-antiinvariant signed measures. We have (compare Section 2 of [8]).

Corollary 2.5: *There is a linear isomorphism Ψ of the kernel of the map $H_b^2(\Gamma, \mathbb{R}) \rightarrow H^2(\hat{S}, \partial\hat{S}, \mathbb{R})$ onto the vector space of all λ -measurable finite flip-antiinvariant Γ -invariant finitely additive signed measures on $S^1 \times S^1 - \Delta$.*

Proof: Every Γ -invariant λ -measurable anti-symmetric cross ratio $[\cdot]$ on S^1 defines a finite finitely additive measurable signed measure $\mu = \Psi([\cdot])$ by assigning to an ordered quadruple (a, b, c, d) of pairwise distinct points in S^1 the value $\mu[a, b] \times [c, d] = [a, b, c, d]$. The map $\Psi : [\cdot] \rightarrow \Psi([\cdot])$ is clearly linear, moreover by Lemma 2.4 its kernel is spanned by the cross ratio $[\cdot]_0$ which defines the Euler class of S viewed as a class in $H^2(\hat{S}, \partial\hat{S}, \mathbb{R})$. As a consequence, the restriction of the map Ψ to the kernel Q of the natural map $H_b^2(\Gamma, \mathbb{R}) \rightarrow H^2(\hat{S}, \partial\hat{S}, \mathbb{R})$ is injective.

Now let μ be any λ -measurable finitely additive Γ -invariant flip anti-invariant signed measure on $S^1 \times S^1 - \Delta$. Define an antisymmetric cross ratio $[\cdot]$ as follows. First, if (a, b, c, d) is ordered and if a, b, c, d are typical points for λ such that $\mu[a, b] \times [c, d]$ is defined then we define $[a, b, c, d] = \mu[a, b] \times [c, d]$. This determines $[a, b, c, d]$ whenever (a, b, c, d) or (b, a, c, d) or (a, b, d, c) is ordered. Choose an arbitrary fixed typical ordered quadruple (a, b, c, d) and define $[a, c, b, d] = 0$. As in the proof of Lemma 2.4, if $b' \in (a, b)$ is arbitrary (and typical) then (c, a, b', b) is ordered and we define $[a, c, b', d] = [a, c, b', b] + [a, c, b, d] = -\mu[c, a] \times [b', b]$. Similarly, for a point $b' \in (a, b)$ and every $a' \in (d, a)$ define $[a', c, b', d] = [a', a, b', d] + [a, c, b', d] = \mu[a', a] \times [b', d] + [a, c, b', d]$. Successively we can define in this way an anti-symmetric measurable Γ -cross ratio on S^1 . As a consequence, the map Ψ is surjective. This shows the corollary. $\square$

It follows from our consideration that there is a direct decomposition $H_b^2(\Gamma, \mathbb{R}) = Q \oplus \ker(\Psi)$ where Q is the kernel of the natural map $H_b^2(\Gamma, \mathbb{R}) \rightarrow H^2(\hat{S}, \partial\hat{S}, \mathbb{R})$ and

Ψ is the map which associates to a λ -measurable anti-symmetric Γ -cross ratio $[\cdot]$ its corresponding finitely additive signed measure on $S^1 \times S^1 - \Delta$.

Remark: Let Γ be a cocompact torsion free lattice in $PSL(2, \mathbb{R})$. Then $S = \mathbb{H}^2/\Gamma$ is a closed surface of genus $g \geq 2$. The *geodesic flow* Φ^t acts on the unit tangent bundle T^1S of S . Call two Hölder continuous functions $f, u : T^1S \rightarrow \mathbb{R}$ *cohomologous* if the integrals of f, u over every periodic orbit for Φ^t coincide. This defines an equivalence relation on the vector space of all Hölder function on T^1S . The *flip* $v \rightarrow -v$ acts on T^1M and induces a map on the space of cohomology classes. We showed in [8] that the space of cohomology classes of *flip anti-invariant* Hölder functions naturally embeds into $H_b^2(\Gamma, \mathbb{R})$. Our above consideration yields another such embedding. Namely, every such function f admits a unique *Gibbs equilibrium state* which defines a Γ -invariant locally finite measure μ on the complement of the diagonal in $S^1 \times S^1$. The function $(a, b, c, d) \rightarrow \mu[a, b] \times [c, d]$ is Hölder continuous. Since f is anti-invariant by assumption, the measure is *not* invariant under the flip unless f is cohomologous to 0. In particular, the signed measure $\tilde{\mu}$ defined by $\tilde{\mu}[a, b] \times [c, d] = \mu[a, b] \times [c, d] - \mu[c, d] \times [a, b]$ is non-trivial, flip anti-invariant and Hölder continuous and hence by Corollary 2.5 it defines a non-trivial class in the kernel of the natural map $H_b^2(\Gamma, \mathbb{R}) \rightarrow H^2(\Gamma, \mathbb{R})$. We do not know whether the signed measures of this form are dense in the space of all finitely additive λ -measurable signed measures equipped with the supremums norm.

3. LARGE GROUPS OF ISOMETRIES OF HYPERBOLIC SPACES

In this section we consider a proper hyperbolic geodesic metric space X . The *Gromov boundary* ∂X of X is defined as follows. For a fixed point $x \in X$, define the *Gromov product* $(y, z)_x$ based at x of two points $y, z \in X$ by

$$(12) \quad (y, z)_x = \frac{1}{2}(d(y, x) + d(z, x) - d(y, z)).$$

Call two sequences $(y_i), (z_j) \subset X$ *equivalent* if $(y_i, z_i)_x \rightarrow \infty$ ($i \rightarrow \infty$). By hyperbolicity of X , this notion of equivalence defines an equivalence relation for the collection of all sequences $(y_i) \subset X$ with the additional property that $(y_i, y_j)_x \rightarrow \infty$ ($i, j \rightarrow \infty$) [3]. The boundary ∂X of X is the set of equivalence classes of this relation.

There is a natural topology on $X \cup \partial X$ which restricts to the given topology on X . With respect to this topology, a sequence $(y_i) \subset X$ converges to $\xi \in \partial X$ if and only if we have $(y_i, y_j)_x \rightarrow \infty$ and the equivalence class of (y_i) defines ξ . Since X is proper by assumption, the space $X \cup \partial X$ is compact and metrizable. Every isometry of X acts naturally on $X \cup \partial X$ as a homeomorphism. Moreover, for every $x \in X$ and every $a \in \partial X$ there is a geodesic ray $\gamma : [0, \infty) \rightarrow X$ with $\gamma(0) = x$ and $\lim_{t \rightarrow \infty} \gamma(t) = a$.

For every proper metric space X the isometry group $\text{Iso}(X)$ of X can be equipped with a natural locally compact σ -compact metrizable topology, the so-called *compact open topology*. With respect to this topology, a sequence $(g_i) \subset \text{Iso}(X)$ converges to some isometry g if and only if $g_i \rightarrow g$ uniformly on compact subsets of X . In this topology, a closed subset $Y \subset \text{Iso}(X)$ is compact if and only if there is

a compact subset K of X such that $gK \cap K \neq \emptyset$ for all $g \in Y$. In particular, the action of $\text{Iso}(X)$ on X is proper. In the sequel we always assume that subgroups of $\text{Iso}(X)$ are equipped with the compact open topology.

The *limit set* Λ of a subgroup G of the isometry group of a proper hyperbolic geodesic metric space X is the set of accumulation points in ∂X of one (and hence every) orbit of the action of G on X . If G is non-compact then its limit set is a compact non-empty G -invariant subset of ∂X . The group G is called *elementary* if its limit set consists of at most two points. In particular, every compact subgroup of $\text{Iso}(X)$ is elementary. If G is non-elementary then its limit set Λ is uncountable without isolated points, and G acts as a group of homeomorphisms on Λ . An element $g \in \text{Iso}(X)$ is called *hyperbolic* if its action on ∂X admits an attracting fixed point $a \in \Lambda$ and a repelling fixed point $b \in \Lambda - \{a\}$ and if it acts on Λ with *north-south-dynamics* with respect to these fixed points.

Assume that $G < \text{Iso}(X)$ is a closed non-elementary subgroup of $\text{Iso}(X)$ with limit set $\Lambda \subset \partial X$. If the group G does not act transitively on the complement of the diagonal in $\Lambda \times \Lambda$ then the kernel of the natural map $H_{cb}^2(G, \mathbb{R}) \rightarrow H_c^2(G, \mathbb{R})$ is infinite dimensional [9]. Extending Lemma 6.1 of [5], we prove Theorem B from the introduction and show that this is not true for groups G which act transitively on the complement of the diagonal in $\Lambda \times \Lambda$.

Proposition 3.1: *Let $G < \text{Iso}(X)$ be a closed non-elementary subgroup with limit set Λ . If G acts transitively on the complement of the diagonal in $\Lambda \times \Lambda$ then the kernel of the natural map $H_{cb}^2(G, \mathbb{R}) \rightarrow H_c^2(G, \mathbb{R})$ is trivial.*

Proof: Let $G < \text{Iso}(X)$ be a closed non-elementary subgroup which acts transitively on the space A of pairs of distinct points in Λ . Then for every fixed point $a \in \Lambda$ the stabilizer G_a of a acts transitively on $\Lambda - \{a\}$.

Following [5], an element in the kernel $H_{cb}^2(G, \mathbb{R}) \rightarrow H_c^2(G, \mathbb{R})$ can be represented by a *continuous quasi-morphism*, i.e. a continuous function $q : G \rightarrow \mathbb{R}$ which satisfies

$$(13) \quad \sup_{g, h \in G} |q(g) + q(h) - q(gh)| < \infty.$$

Such a continuous quasi-morphism is bounded on every *compact* subset of G and is bounded on each fixed conjugacy class in G . The quasi-morphism q defines a non-trivial element of $H_{cb}^2(G, \mathbb{R})$ only if q is unbounded.

Since G is non-elementary by assumption, G contains a hyperbolic element $g \in G$. Let $a \in \Lambda$ be the attracting fixed point of g and let $b \in \Lambda - \{a\}$ be the repelling fixed point. Then g preserves the set of geodesics connecting b to a ; note that such a geodesic exists since X is proper. For $x, y \in X$ and a geodesic ray $\gamma : [0, \infty) \rightarrow X$ connecting x to a write $\beta(y, \gamma) = \limsup_{t \rightarrow \infty} (d(y, \gamma(t)) - t)$ and define the *Busemann function*

$$(14) \quad \beta_a(y, x) = \sup\{\beta(y, \gamma) \mid \gamma \text{ is a geodesic ray connecting } x \text{ to } a\}.$$

Then we have $|\beta_a(y, x)| \leq d(y, x)$, moreover there is a number $c > 0$ only depending on the hyperbolicity constant for X such that

$$(15) \quad |\beta_a(\cdot, y) - \beta_a(\cdot, x) + \beta_a(y, x)| \leq c$$

for all $x, y \in X$ (Proposition 8.2 of [7]). Let $\gamma : [0, \infty) \rightarrow X$ be a geodesic ray connecting x to a . By hyperbolicity, for every $t > 0$ every geodesic ray connecting x to a passes through a neighborhood of $\gamma(t)$ of uniformly bounded diameter and therefore via possibly enlarging c we may assume that

$$(16) \quad |\beta_a(\gamma(t), x) + d(\gamma(t), x)| \leq c$$

independent of x, γ, t . Define the *horosphere* at a through x to be the set $H_a(x) = \beta_a(\cdot, x)^{-1}(0)$. For all $x, y \in X$ the distance between the horospheres $H_a(x), H_a(y)$ is not smaller than $|\beta_a(x, y)| - c$. Moreover, the estimates (15), (16) show that for all x, y with $\beta_a(x, y) \leq 0$ the distance between y and $H_a(x)$ is not bigger than $|\beta_a(x, y)| + 2c$.

Let $W(a, b) \subset X$ be the closed non-empty subset of all points in X which lie on a geodesic connecting b to a . The isometry $g \in G$ is hyperbolic with fixed points $a, b \in \Lambda$ and therefore it preserves $W(a, b)$. If we denote by Γ the infinite cyclic subgroup of G generated by g then $W(a, b)/\Gamma$ is compact. As a consequence, there is a number $\nu > 0$ and for every $x \in W(a, b)$ and every $t \in \mathbb{R}$ there is a number $k(t) \in \mathbb{Z}$ with $|\beta_a(g^{k(t)}x, x) - t| < \nu$. It follows from this and (16) that for every $y \in X$ there is some $k = k(y) \in \mathbb{Z}$ such that the distance between $g^k x$ and the horosphere $H_a(y)$ is at most $\delta_0 = \nu + 2c$.

By assumption, the stabilizer $G_a < G$ of the point $a \in \Lambda$ acts transitively on $\Lambda - \{a\}$. Thus there is for every $\zeta \in \Lambda - \{a\}$ an element $h_\zeta \in G_a$ with $h_\zeta(b) = \zeta$. Let $x \in W(a, b)$; since $h_\zeta \in G_a$ we have $h_\zeta^{-1}(H_a(x)) = H_a(h_\zeta^{-1}(x))$. By our above consideration, there is a number $\ell \in \mathbb{Z}$ such that the distance between $g^\ell(x)$ and $H_a(h_\zeta^{-1}(x))$ is at most δ_0 and therefore the distance between $h_\zeta \circ g^\ell(x)$ and $H_a(x)$ is at most δ_0 . Hence via replacing h_ζ by $h_\zeta \circ g^\ell \in G_a$ we may assume that the distance between $h_\zeta(x)$ and $H_a(x)$ is at most δ_0 . In particular, we have $|\beta_a(\cdot, x) - \beta_a(\cdot, h_\zeta(x))| \leq \delta_0 + c$. Now h maps the geodesic γ connecting x to a to a geodesic $h\gamma$ connecting $h_\zeta x$ to a and hence by the definition of the Busemann functions, for every $t \in \mathbb{R}$ the distance between $h_\zeta(\gamma(t))$ and $H_a(\gamma(t))$ is bounded from above by $\delta_0 + 3c$. Since Γ preserves the set of geodesics connecting b to a we conclude that there is a universal constant $\delta_1 > 0$ such that for every $\ell \geq 0$ the distance between $h_\zeta(g^\ell x)$ and $H_a(g^\ell(x))$ is at most δ_1 .

For $x \in X$ denote by $N_{a,x} \subset G_a$ the set of all elements $h \in G_a$ with the property that the distance between $H_a(x)$ and hx is at most δ_0 . Our above consideration shows that for every $x \in W(a, b)$ and every $\zeta \in \Lambda - \{a\}$ there is some $h_\zeta \in N_{a,x}$ which maps b to ζ . For every $h \in N_{a,x}$ the sequence $(g^{-\ell} \circ h \circ g^\ell)_\ell$ is contained in G_a and by the above consideration, it maps x into the δ_1 -neighborhood of $H_a(x)$. Moreover, the sequence $(g^{-\ell} \circ h \circ g^\ell(b))_\ell \subset \Lambda$ converges as $\ell \rightarrow \infty$ to b . This means that there is a number $\delta_2 > \delta_1$ such that for sufficiently large ℓ the element $g^{-\ell} \circ h \circ g^\ell \in G_a$ maps the point x into the closed δ_2 -neighborhood B_x of x . The group Γ acts on $W(a, b)$ cocompactly and hence if $C \subset W(a, b)$ is a compact fundamental domain for this action then $B = \cup_{x \in C} B_x$ is compact, moreover for

every $x \in W(a, b)$, every element $h \in N_{a,x}$ is conjugate in G to an element in the compact subset $K = \{u \in G \mid uB \cap B \neq \emptyset\}$ of G . As a consequence, the restriction to $N_{a,b} = \cup_{x \in W(a,b)} N_{a,x}$ of any continuous quasi-morphism q on G is uniformly bounded. By our assumption on G the sets $N_{a,b}$ ($(a, b) \in A$) are pairwise conjugate in G and hence q is uniformly bounded on $\cup_{(a,b) \in A} N_{a,b} = N$.

Next we show that the restriction of a quasi-morphism q to the subgroup $G_{a,b}$ of G_a which stabilizes both points $a, b \in \Lambda$ is bounded. For this consider an arbitrary element $u \in G_{a,b}$. Let $x \in W(a, b)$ and let $\beta_a(ux, x) = \tau$; we may assume that $u \notin N_{a,b}$ and hence after possibly exchanging b and a that $\tau < 0$. Since u preserves $W(a, b)$, by (16) we have $|d(ux, x) + \tau| \leq \tilde{c}$ where $\tilde{c} > 0$ is a universal constant. Choose some $h \in G$ with $h(a, b) = (b, a)$; then h preserves the set $W(a, b)$. Let $g \in G_{a,b}$ be the fixed hyperbolic element as before. By our above consideration, via composing h with g^ℓ for a number $\ell \in \mathbb{Z}$ depending on h we may assume that $d(x, hx) \leq \delta_2$ for a universal constant $\delta_2 > 0$. We have $hux \in W(a, b)$ and $|\beta_b(hux, x) - \beta_a(ux, x)| \leq |\beta_b(hux, hx) - \beta_a(ux, x)| + d(hx, x) \leq \delta_2$. On the other hand, from another application of the estimate (16) we obtain the existence of a constant $\delta_3 > 0$ such that $|\beta_b(z, y) - \beta_a(y, z)| \leq \delta_3$ for all $y, z \in W(a, b)$. But this just means that the distance between hux and $u^{-1}x$ is uniformly bounded and hence the distance between $uhux$ and x is uniformly bounded as well. As a consequence, the element uhu is contained in a fixed compact subset of G . As before, we conclude from this that $|q(u)|$ is bounded from above by a universal constant not depending on u and hence the restriction of q to $G_{a,b}$ is uniformly bounded.

Now let $h \in G$ be arbitrary and assume that $ha = x, hb = y$. We showed above that there are $h_y \in N \cap G_x, h_x \in N \cap G_b$ with $h_y(y) = b$ and $h_x(x) = a$; then $h' = h_x h_y h \in G_{a,b}$ and $|q(h') - q(h)|$ is uniformly bounded. As a consequence, q is bounded and hence it defines the trivial bounded cohomology class. This completes the proof of the proposition. $\square$

From now on we assume that X is of *bounded growth* which means that there is a number $b > 1$ such that for every $R > 1$, every metric ball of radius R contains at most be^{bR} disjoint metric balls of radius 1. Let $G < \text{Iso}(X)$ be a closed non-elementary group with limit set Λ . Let (B, μ_0) be a strong boundary for G . By Lemma 2.2 of [9] there is a G -equivariant map $\varphi : B \rightarrow \Lambda$. The image μ of μ_0 under the map φ is a measure on Λ whose measure class is invariant under the action of G and such that the diagonal action of G on $\Lambda \times \Lambda$ is ergodic with respect to the class of $\mu \times \mu$. Since X is of bounded growth by assumption, the stabilizer in $\text{Iso}(X)$ of each point in ∂X is amenable and therefore the measure space (Λ, μ) is a strong boundary for G (compare the discussion in [1, 2, 11]). In particular, Corollary 2.3 implies that $H_{cb}^2(G, \mathbb{R})$ is isomorphic to the space of μ -measurable anti-symmetric G -cross ratios on Λ .

Now assume that G acts transitively on the complement of the diagonal in $\Lambda \times \Lambda$. The stabilizer $G_{a,b}$ of (a, b) acts on $\Lambda - \{a, b\}$ as a group of homeomorphisms. Moreover, there is some $g \in G$ which maps (a, b) to (b, a) . Define the group G to be *directed* if there is a $G_{a,b}$ -invariant subset $U(a, b)$ of $\Lambda - \{a, b\}$ of positive μ -mass such that $gU(a, b) \cap U(a, b) = \emptyset$. We have.

Lemma 3.2: *Let $G < \text{Iso}(X)$ be a closed subgroup with limit set Λ . Assume that G acts transitively on the complement A of the diagonal in $\Lambda \times \Lambda$; if $H_{cb}^2(G, \mathbb{R}) \neq \{0\}$ then G is directed.*

Proof: Let $G < \text{Iso}(X)$ be a closed subgroup with limit set Λ which acts transitively on the complement A of the diagonal in $\Lambda \times \Lambda$. If $H_{cb}^2(G, \mathbb{R}) \neq 0$ then there is some G -invariant measurable nontrivial alternating function φ on the space of triples of pairwise distinct points in Λ which satisfies the cocycle identity (4).

Let φ be any G -invariant μ -measurable alternating function on the space of triples of pairwise distinct points in Λ . Let $(a, b) \in A$; since the action of G on A is transitive, there is a set $U \subset \Lambda$ of positive μ -mass such that $\varphi(a, b, u) > 0$ for every $u \in U$. If $v = hu$ for some $h \in G_{a,b}$ then $\varphi(a, b, u) = \varphi(a, b, v)$. Let $g \in G$ be such that $g(a, b) = (b, a)$. Then $g^{-1}G_{a,b}g = G_{a,b}$ and if G is not directed then for every $G_{a,b}$ -invariant measurable subset U of $\Lambda - \{a, b\}$ with $\mu(U) > 0$ we have $\mu(U \cap gU) > 0$. Now for every $c \in \mathbb{R}$ and every $\epsilon > 0$ the set $U_{c,\epsilon} = \{u \mid \varphi(a, b, u) \in (c - \epsilon, c + \epsilon)\}$ is $G_{a,b}$ -invariant. By assumption, if $\mu(U_{c,\epsilon}) > 0$ then we have $\mu(gU_{c,\epsilon} \cap U_{c,\epsilon}) > 0$ as well and therefore there is some $u \in U_{c,\epsilon}$ with $|\varphi(a, b, gu) - c| < \epsilon$. Since φ is alternating we obtain that $\varphi(a, b, gu) = -\varphi(b, a, gu) = -\varphi(ga, gb, gu) = -\varphi(a, b, u) \in (-c - \epsilon, -c + \epsilon)$ and consequently $|c| < 2\epsilon$. On the other hand, for μ -almost every $u \in \Lambda$ and every $\epsilon > 0$ we have $\mu(U_{\varphi(a,b,u),\epsilon}) > 0$. From this we deduce that φ vanishes almost everywhere and hence $H_{cb}^2(G, \mathbb{R}) = \{0\}$. $\square$

In the case that G is a simple Lie group of non-compact type and rank 1 we have $H_{cb}^2(G, \mathbb{R}) \neq \{0\}$ if and only if $G = SU(n, 1)$ for some $n \geq 1$. That this condition is necessary is immediate from Lemma 3.2 and the following observation.

Lemma 3.3: *A simple Lie group G of non-compact type and rank one is directed only if $G = SU(n, 1)$ for some $n \geq 1$.*

Proof: Let G be a simple Lie group of non-compact type and rank one. Then G is the isometry group of a symmetric space X of non-compact type and negative curvature with ideal boundary ∂X . The action of G on ∂X preserves the measure class of the Lebesgue measure λ and $(\partial X, \lambda)$ is a strong boundary for G . The group G acts transitively on the space A of pairs of distinct points in ∂X . If $G = SO(n, 1)$ for some $n \geq 3$ then G acts transitively on the space of triples of pairwise distinct points in ∂X and hence by the above, G is not directed.

Now let $G = Sp(n, 1)$ for some $n \geq 1$. Then for every pair $(a, b) \in A$ there is a unique totally geodesic embedded quaternionic line $L \subset X$ of constant curvature -4 whose boundary $\partial L = S^3 \subset \partial X$ contains a and b . The stabilizer $G_{a,b}$ of (a, b) is contained in the stabilizer G_L of L in G which is conjugate to the quotient of the group $Sp(1, 1) \times Sp(n-1) < Sp(n, 1)$ by its center. Moreover, G_L acts transitively on the space of triples of pairwise distinct points in ∂L . In particular, for every $u \in \partial L$ and every $g \in G$ with $g(a, b) = (b, a)$ the $G_{a,b}$ -orbit of u coincides with the $G_{a,b}$ -orbit of gu . More generally, this is true for every point $u \in \partial X - \{a, b\}$. Namely, let $P : \partial X \rightarrow L$ be the shortest distance projection. The subgroup $\tilde{G}_{Pu}$ of G_L which stabilizes Pu is the quotient of the group $Sp(1) \times Sp(n-1)$ by its

center. The factor subgroup $Sp(n-1)$ acts transitively on $P^{-1}(Pu)$ while the orbit of Pu under the group $G_{a,b}$ consists of the set of all points in L whose distance to the geodesic connecting b to a coincides with the distance of Pu . As above we conclude that for every $u \in \partial X$ and every $g \in G$ with $g(a,b) = (b,a)$ the $G_{a,b}$ -orbit of u coincides with the $G_{a,b}$ -orbit of gu . In other words, G is not directed. In the same way we obtain that the exceptional Lie group F_{20}^4 is not directed as well. $\square$

Note that for every $n \geq 1$ the group $SU(n,1)$ admits a non-trivial continuous second bounded cohomology class $\alpha \in H_{cb}^2(SU(n,1), \mathbb{R})$ induced by the Kähler form ω of the complex hyperbolic space $X = SU(n,1)/S(U(n)U(1))$, and this class generates $H_{cb}^2(SU(n,1), \mathbb{R})$. The anti-symmetric cross ratio $[\cdot]$ on ∂X defining the class α is a continuous function on ∂X with values in $[-\pi, \pi]$. We have $|[a, b, c, d]| = \pi$ if and only if (a, c, b, d) or (c, a, d, b) is an ordered quadruple of points in the boundary of a complex line in X (for a discussion of this fact, references and applications to rigidity, see the recent preprint of Burger and Iozzi [4]).

4. RIGIDITY OF COCYCLES

In this section we consider as before a proper hyperbolic geodesic metric space (X, d) . We equip the isometry group $\text{Iso}(X)$ with the compact open topology; with respect to this topology, $\text{Iso}(X)$ is a locally compact σ -compact topological group. Let S be a standard Borel space and let μ be a Borel probability measure on S . Assume that G is a locally compact σ -compact topological group which admits a measure preserving ergodic action on a standard probability space (S, μ) . This action then defines a natural continuous unitary representation of G into the Hilbert space $L^2(S, \mu)$ of square integrable functions on S . Let $\alpha : G \times S \rightarrow \text{Iso}(X)$ be a cocycle, i.e. $\alpha : G \times S \rightarrow \text{Iso}(X)$ is a Borel map which satisfies $\alpha(gh, x) = \alpha(g, hx)\alpha(h, x)$ for all $g, h \in G$ and μ -almost every $x \in S$. The cocycle is *cohomologous* to a cocycle $\beta : G \times S \rightarrow \text{Iso}(X)$ if there is a measurable function $\psi : S \rightarrow \text{Iso}(X)$ such that $\psi(gx)\alpha(g, x) = \beta(g, x)\psi(x)$ for all $g \in G$ and almost all $x \in S$. We use Theorem 4.1 of [9] to show (compare [14], [12]).

Lemma 4.1: *Let G be a locally compact σ -compact group and let (S, μ) be an ergodic G -probability space. Let X be a proper hyperbolic geodesic metric space and let $\alpha : G \times S \rightarrow \text{Iso}(X)$ be a cocycle. Let $H < \text{Iso}(X)$ be a closed subgroup with limit set Λ and assume that $\alpha(G \times S) \subset H$ but that α is not cohomologous to a cocycle into a proper subgroup of H . If $H_{cb}^2(G, L^2(S, \mu))$ is finite dimensional then either H is elementary or H acts transitively on the complement of the diagonal in $\Lambda \times \Lambda$.*

Proof: Let G be a locally compact σ -compact group which admits an ergodic measure preserving action on a standard Borel probability space (S, μ) . Let $\alpha : G \times S \rightarrow \text{Iso}(X)$ be a cocycle and assume that $\alpha(G \times S)$ is contained in a closed subgroup H of $\text{Iso}(X)$ and that α is not equivalent to a cocycle with values in a proper closed subgroup of H . Assume moreover that H is not elementary and does not act transitively on the complement of the diagonal in $\Lambda \times \Lambda$. By Theorem 4.1 of [9], $H_{cb}^2(H, \mathbb{R})$ is infinite dimensional. More precisely, if $T \subset \Lambda^3$ is the space of

triples of pairwise distinct points in Λ then there is an infinite dimensional vector space of *continuous* H -invariant cocycles on T , i.e. continuous H -invariant maps $T \rightarrow \mathbb{R}$ which satisfy the cocycle equation (5).

Let (B, ν) be a strong boundary for G . By Lemma 2.2 of [9] there is an α -equivariant measurable map $\varphi : B \times S \rightarrow \Lambda$. This means that for almost every $(w, \sigma) \in B \times S$ and every $g \in G$ we have $\varphi(gw, g\sigma) = \alpha(g, \sigma)\varphi(w, \sigma)$. We claim that the image of φ is dense in Λ . Namely, otherwise there is a proper closed subset A of Λ which contains the image of φ . Then for every $g \in G$ and almost every $u \in S$ the element $\alpha(g, u) \in H$ stabilizes A , and α is cohomologous to a cocycle with values in the intersection of H with the stabilizer of A . Since A is a closed proper subset of Λ , the stabilizer of A intersects H in a closed subgroup H' of infinite index which contradicts our assumption on α .

We now follow Monod and Shalom [14]. Namely, choose a continuous H -invariant bounded cocycle $\omega : T \rightarrow \mathbb{R}$. Define a measurable map $\eta(\omega)$ from the space $\tilde{B}$ of triples of pairwise distinct points in B into $L^\infty(S, \mu)$ by $\eta(x_1, x_2, x_3)(z) = \omega(\varphi(x_1, z), \varphi(x_2, z), \varphi(x_3, z))$; then η can be viewed as a measurable G -equivariant bounded cocycle with values in the Hilbert space of square integrable function on S . Since ω is continuous and $\varphi(B \times S)$ is dense in Λ , this cocycle does not vanish identically. As a consequence, the space of G -equivariant measurable bounded cocycles $\tilde{B} \rightarrow L^2(S, \mu)$ is infinite dimensional. On the other hand, B is a strong boundary for G and therefore $H_b^2(G, L^2(S, \mu))$ is isomorphic to the space of measurable G -equivariant bounded cocycles $\tilde{B} \rightarrow L^2(S, \mu)$ [13]. In other words, $H_{cb}^2(G, L^2(S, \mu))$ is infinite dimensional. This shows the lemma. $\square$

Now consider a semi-simple Lie group G with finite center and no compact factors. Let $\Gamma < G$ be an irreducible lattice in G , i.e. Γ is a discrete subgroup of G such that the volume of G/Γ is finite and that moreover if $G = G_1 \times G_2$ is a non-trivial product then the projection of Γ to each factor group G_i ($i = 1, 2$) is dense. The group G acts on G/Γ preserving the projection of the Lebesgue measure λ on G .

Let (S, μ) be a standard probability space with a measure preserving action of Γ . Then Γ admits a measure preserving action on the product space $G \times S$. Since $\Gamma < G$ is a lattice, the quotient space $(G \times S)/\Gamma$ can be viewed as a bundle over G/Γ with fibre S . If $\Omega \subset G$ is a finite measure Borel fundamental domain for the action of Γ on G then $\Omega \times S \subset G \times S$ is a finite measure Borel fundamental domain for the action of Γ on $G \times S$ and up to normalization, the product measure $\lambda \times \mu$ projects to a finite measure ν on $(G \times S)/\Gamma$. The action of G on $G \times S$ by left translation commutes with the action of Γ and hence it projects to an action on $(G \times S)/\Gamma$ preserving the measure ν (see p.75 of [16]). We have.

Lemma 4.2: *Let Γ be an irreducible lattice in a product $G = G_1 \times G_2$ of two semi-simple non-compact Lie groups G_1, G_2 . If the action of Γ on (S, μ) is mixing then the induced action of G_1 on $((G \times S)/\Gamma, \nu)$ is ergodic.*

Proof: Let Γ be an irreducible lattice in a product $G = G_1 \times G_2$ of two semi-simple non-compact Lie groups G_1, G_2 with finite center. Let (S, μ) be a mixing

Γ -probability space. The induced action of G on $(G \times S)/\Gamma$ preserves the measure ν and restricts to an action of G_1 . Since the measure ν is finite, this action is ergodic if and only if every G_1 -invariant function $f \in L^2((G \times S)/\Gamma, \nu)$ is constant.

We argue by contradiction and we assume that there is a G_1 -invariant function $f \in L^2((G \times S)/\Gamma, \nu)$ with $\int \|f\|^2 d\nu = 1$ and zero mean $\int f d\nu = 0$. Let $\tilde{f}$ be the lift of f to a Γ -invariant function on $G \times S$; then $\tilde{f}$ is locally square integrable with respect to the product measure $\lambda \times \mu$. For $x \in G$ define a function $\varphi(x)$ on S by $\varphi(x)(y) = \tilde{f}(x, y)$; by Fubini's theorem we have $\varphi(x) \in L^2(S, \mu)$ for λ -almost every x , moreover the function $x \rightarrow \int_S \varphi(x) d\mu$ is measurable with respect to λ . We claim that $\int_S \varphi(x) d\mu = 0$ and $\int_S |\varphi(x)|^2 d\mu = 1$ for λ -almost every $x \in G$.

For this recall that by definition of the action of G_1 , the functions $x \rightarrow \int_S \varphi(x) d\mu$ and $x \rightarrow \int_S |\varphi(x)|^2 d\mu$ are invariant under the action of G_1 on G/Γ since the action of Γ on (S, μ) is measure preserving. By Moore's ergodicity theorem the action of G_1 on G/Γ is ergodic [16] and hence these functions are constant almost everywhere. If $\Omega \subset G$ is a Borel-fundamental domain for the action of Γ on G and if the Lebesgue measure λ on G is normalized in such a way that $\lambda(\Omega) = 1$ then $\int_\Omega \int_S \varphi(x) d\mu d\lambda = 0$, $\int_\Omega \int_S |\varphi(x)|^2 d\mu d\lambda = 1$ and hence these constants equal 0, 1. In particular, if we denote by $\mathcal{V}$ the orthogonal complement in $L^2(S, \mu)$ of the constant functions then the assignment $x \rightarrow \varphi(x)$ is a $\mathcal{V}$ -valued measurable function on G/Γ .

Let $L^{[1]}(G, \mathcal{V})^\Gamma$ be the space of all Γ -invariant measurable maps $h : G \rightarrow \mathcal{V}$ for which the function $\|h\|^2$ is locally integrable with respect to the Lebesgue measure λ on G . Note that $L^{[1]}(G, \mathcal{V})^\Gamma$ naturally has the structure of a separable Banach space. The group G_1 acts isometrically on $L^{[1]}(G, \mathcal{V})^\Gamma$ by left translation. By our assumption, there is a non-zero G_1 -invariant vector $\varphi \in L^{[1]}(G, \mathcal{V})^\Gamma$. Then $\varphi : G \rightarrow \mathcal{V}$ is measurable and essentially bounded. Thus the restriction of φ to suitable compact sets of large measure is continuous; in particular, we can find a compact set $K = K_1 \times K_2 \subset G_1 \times G_2$ of positive measure, Borel sets $D, E \subset S$ of positive measure and a number $\epsilon > 0$ such that the following is satisfied.

- (1) $\varphi(z)(u) \geq \epsilon$ for every $z \in K, u \in D$.
- (2) $\varphi(z)(v) \leq 0$ for every $z \in K, v \in E$.

Since φ is invariant under the left action of G_1 , for any $z_0 \in K_1, z_2 \in K_2$ and $z_1 \in G_1, u \in S$ we have $\varphi(z_1, z_2)(u) = \varphi(z_0, z_2)(u)$. On the other hand, φ is also invariant under the right action of Γ and hence $\varphi(z_1, z_2)(u) = \varphi((z_1, z_2)\eta)(\eta u)$ for every $\eta \in \Gamma$. In particular, if $(z_1, z_2)\eta \in K$ then we deduce that $\mu(\eta D \cap E) = 0$.

The right action of Γ on $G_1 \backslash G$ is ergodic and therefore there are infinitely many elements $\eta \in \Gamma$ such that the measure of $(G_1 \times K_2)\eta \cap K$ is positive. Moreover, the action of Γ on (S, μ) is mixing by assumption and hence for every $\eta \in \Gamma$ which is sufficiently far away from the identity the image of D under η intersects E in a set of positive measure. In particular, there is some $\eta \in \Gamma$ with $\lambda((G_1 \times K_2)\eta \cap K) > 0$ and $\mu(\eta D \cap E) > 0$ which contradicts the above consideration. Thus the vector space of G_1 -invariant maps in $L^{[1]}(G, \mathcal{V})^\Gamma$ is trivial and the action of G_1 on $(G \times S)/\Gamma$ is ergodic. $\square$

The following corollary completes the proof of Theorem A from the introduction and follows as in [14] from Lemma 4.2, the rigidity results for bounded cohomology of Burger and Monod [5, 6] and the work of Zimmer [16].

Corollary 4.3: *Let G be a semi-simple Lie group with finite center, no compact factors and of rank at least 2. Let $\Gamma < G$ be an irreducible lattice and let (S, μ) be a mixing Γ -space. Let X be a proper hyperbolic geodesic metric space and let $\alpha : \Gamma \times S \rightarrow \text{Iso}(X)$ be a cocycle; then either α is cohomologous to a cocycle into an elementary subgroup of $\text{Iso}(X)$ or there is a closed subgroup H of $\text{Iso}(X)$ which is a compact extension of a simple Lie group L of rank one and there is a surjective homomorphism $G \rightarrow L$.*

Proof: Let G be a semi-simple Lie group of non-compact type with finite center and of rank at least 2 and let Γ be an irreducible lattice in G . Let (S, μ) be a mixing Γ -space with invariant Borel probability measure μ . Let $\alpha : \Gamma \times S \rightarrow \text{Iso}(X)$ be a cocycle into the isometry group of a proper hyperbolic geodesic metric space X . Let $H < \text{Iso}(X)$ be a closed subgroup such that α is cohomologous to an H -valued cocycle but not to a cocycle with values in a proper subgroup of H . We may assume that $\alpha(\Gamma \times S) \subset H$. Assume that H is not elementary, with limit set Λ . By Theorem A in [9], there is a continuous $L^2(H)$ -valued bounded cocycle $\varphi : \Lambda^3 \rightarrow L^2(H)$.

Let $\Omega \subset G$ be a Borel fundamental domain for the action of Γ on G . Let ν be the G -invariant Borel probability measure on $(G \times S)/\Gamma \sim \Omega \times S$. We obtain a ν -measurable function $\beta : G \times (G \times S)/\Gamma \rightarrow H$ as follows. For $z \in \Omega$ and $g \in G$ let $\eta(g, z) \in \Gamma$ be the unique element such that $gz \in \eta(g, z)\Omega$ and define $\beta(g, (z, \sigma)) = \alpha(\eta(g, z), \sigma)$. By construction, η satisfies the cocycle equation for the action of G on $(G \times S)/\Gamma$. Let (B, μ) be a strong boundary for G ; we may assume that B is also a strong boundary for Γ . By Lemma 4.2 the action of G on $(G \times S)/\Gamma$ is ergodic and hence Lemma 2.2 of [9] show that we can find a measurable β -equivariant Fürstenberg map $\psi_0 : (G \times S)/\Gamma \times B \rightarrow \Lambda$ and hence a β -equivariant map $\psi : (G \times S)/\Gamma \times B^3 \rightarrow \Lambda^3$ defined by $\psi(x, a, b, c) = (\psi_0(x, a), \psi_0(x, b), \psi_0(x, c))$ ($x \in (G \times S)/\Gamma$). Let $L^{[2]}((G \times S)/\Gamma, L^2(H))$ be the space of all measurable maps $(G \times S)/\Gamma \rightarrow L^2(H)$ with the additional property that for each such map φ the function $x \rightarrow \|\varphi\|$ is square integrable on $(G \times S)/\Gamma$. Then $L^{[2]}((G \times S)/\Gamma, L^2(H))$ has a natural structure of a separable Hilbert space, and the group G acts on $L^{[2]}((G \times S)/\Gamma, L^2(H))$ as a group of isometries. In other words, $L^{[2]}((G \times S)/\Gamma, L^2(H))$ is a Hilbert module for G and the cocycle φ can be composed with the map ψ to a β -invariant measurable bounded map $B^3 \rightarrow L^{[2]}((G \times S)/\Gamma, L^2(H))^3$. Since B is a strong boundary for G we conclude that this map defines a nontrivial cohomology class in $H_{cb}^2(G, L^{[2]}((G \times S)/\Gamma, L^2(H)))$.

Now if G is simple then the results of Monod and Shalom [14] show that there is a β -equivariant map $(G \times S)/\Gamma \rightarrow L^2(H)$. Since the action of G on $(G \times S)/\Gamma$ is ergodic, by the cocycle reduction lemma of Zimmer [16] the cocycle β and hence α is cohomologous to a cocycle into a compact subgroup of H which is a contradiction. On the other hand, if $G = G_1 \times G_2$ for semi-simple Lie groups G_1, G_2 with finite center and without compact factors, then the results of Burger and Monod [5, 6] show that via possibly exchanging G_1 and G_2 we may assume that there is an

equivariant map $(G \times S)/\Gamma \rightarrow L^2(H)$ for the restriction of β to $G_1 \times (G \times S)/\Gamma$, viewed as a cocycle for G_1 . By Lemma 4.2 the action of G_1 on $(G \times S)/\Gamma$ is ergodic and therefore the cocycle reduction lemma of Zimmer [16] shows that the restriction of β to G_1 is equivalent to a cocycle into a compact subgroup of H . We now follow the proof of Theorem 1.2 of [14] and find a minimal such compact subgroup K of H . The cocycle β and hence α is cohomologous to a cocycle into the normalizer of K in H which then coincides with H by our assumption of H . Moreover, there is a continuous homomorphism of G onto H/K . Since G is connected, the image of G under this homomorphism is connected as well and hence by Theorem A of [9], H/K is a simple Lie group of rank one. This completes the proof of the corollary. $\square$

REFERENCES

- [1] S. Adams, *Boundary amenability for word hyperbolic groups and an application to smooth dynamics of simple groups*, Topology 33 (1994), 765–783.
- [2] S. Adams, *Reduction of cocycles with hyperbolic targets*, Erg. Th. & Dyn. Sys. 16 (1996), 1111–1145.
- [3] M. Bridson, A. Haefliger, *Metric spaces of non-positive curvature*, Springer, Berlin Heidelberg 1999.
- [4] M. Burger, A. Iozzi, *Bounded cohomology theory and applications to deformation rigidity in complex hyperbolic geometry*, preprint 2005.
- [5] M. Burger, N. Monod, *Bounded cohomology of lattices in higher rank Lie groups*, J. Eur. Math. Soc. 1 (1999), 199–235.
- [6] M. Burger, N. Monod, *Continuous bounded cohomology and applications to rigidity theory*, Geom. Funct. Anal. 12 (2002), 219–280.
- [7] E. Ghys, P. de la Harpe, *Sur les groupes hyperboliques d’après Mikhael Gromov*, Birkhäuser, Boston 1990.
- [8] U. Hamenstädt, *Bounded cohomology and isometry groups of hyperbolic spaces*, arXiv:math.GR/0507097.
- [9] U. Hamenstädt, *Isometry groups of proper hyperbolic spaces*, arXiv:math.GR/0507608.
- [10] N. V. Ivanov, *Foundations of the theory of bounded cohomology*, J. Soviet Math. 37 (1987), 1090–1115.
- [11] V. Kaimanovich, *Double ergodicity of the Poisson boundary and applications to bounded cohomology*, Geom. Funct. Anal. 13 (2003), 852–861.
- [12] I. Mineyev, N. Monod, Y. Shalom, *Ideal bicomings for hyperbolic groups and applications*, Topology 43 (2004), 1319–1344.
- [13] N. Monod, *Continuous bounded cohomology of locally compact groups*, Lecture Notes in Math. 1758, Springer 2001.
- [14] N. Monod, Y. Shalom, *Cocycle superrigidity and bounded cohomology for negatively curved spaces*, J. Diff. Geom. 67 (2004), 395–456.
- [15] N. Monod, Y. Shalom, *Orbit equivalence rigidity and bounded cohomology*, to appear in Ann. Math.
- [16] R. Zimmer, *Ergodic theory and semisimple groups*, Birkhäuser, Boston 1984.

MATHEMATISCHES INSTITUT DER UNIVERSITÄT BONN
 BERLINGSTRASSE 1
 D-53115 BONN

e-mail: ursula@math.uni-bonn.de